Florent Jacquemard

Reachability and confluence are undecidable for flat term rewriting systems

Research Report LSV–03–6, Mar. 2003

Laboratoire Spécification et Vérification

Ecole Normale Supérieure de Cachan
61, avenue du Président Wilson
94235 Cachan Cedex France

CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

http://www.lsv.ens-cachan.fr/Publis/
Research Report LSV-03-6, Lab. Spécification et Vérification, CNRS & ENS de Cachan, France, Mar. 2003

Final version published in *Information Processing Letters* 87(5):265–270, 2003.

Reachability and confluence are undecidable for flat term rewriting systems

Florent Jacquemard

INRIA Futurs & LSV/CNRS UMR 8643, ENS de Cachan,
61, avenue du président-Wilson
94235 Cachan Cedex, France
Tel: +33-1 47 40 75 44 Fax: +33-1 47 40 24 64

Abstract

Ground reachability, ground joinability and confluence are shown undecidable for flat term rewriting systems, i.e. systems in which all left and right members of rule have depth at most one.

Key words: Theory of computation, Term rewriting.

Introduction

The confluence of a term rewriting systems (TRS) guaranties that every term has at most one normal form. This property is undecidable in general, and has been shown decidable for ground TRS in [1,2]. The main result of [1] also implies the decidability of the reachability and joinability problems for ground TRS.

More recently, the confluence has been shown solvable in polynomial time for several classes of TRS, every class embedding the previous one: some restricted ground TRS in [3], ground TRS in [4], shallow (variables occur at depth at most 1 in rewrite rules) and rule linear (in every rewrite rule, every variable occurs at most once) TRS in [5], and shallow, linear (in every left or right member of rewrite rule, every variable occurs at most once) TRS in [6]. The polynomial time complexity result of [4] is also valid for the decision of reachability and joinability, which was already shown in [1].

Email address: florent.jacquemard@lsv.ens-cachan.fr (Florent Jacquemard).

Reachability, joinability and confluence are undecidable for linear (non-shallow) TRS [7], but it was not known whether we can relax the linearity assumptions on variables of the systems of [5,6], keeping these properties decidable [1]. We answer here by the negative, showing, with a reduction of the Post Correspondence Problem, that the problems of ground reachability, ground joinability and confluence are undecidable for flat TRS (every terms in rewrite rules have depth at most 1) with non linear variables. The proof for ground reachability uses the same *colored* techniques as an older proof of undecidability of rigid reachability [8], though this latter result could not be reused directly in this context.

1 Preliminaries

Given a signature Σ, and a set of variable symbols $\mathcal{X}$, we note $\mathcal{T}(\Sigma, \mathcal{X})$ the set of terms build with symbols of Σ and $\mathcal{X}$ and $\mathcal{T}(\Sigma)$ its subset of *ground terms*. The set of function symbols of Σ of arity i is denoted Σ_i.

A *term rewriting system* (TRS) on Σ is defined as a finite set of rewrite rules denoted $\ell \to r$ with $\ell, r \in \mathcal{T}(\Sigma, \mathcal{X})$. We note $\xrightarrow[R]{}$ the rewrite relation (on terms of $\mathcal{T}(\Sigma, \mathcal{X})$) defined by the TRS R, and $\xrightarrow[R]{*}$ the reflexive and transitive of this relation.

Definition 1 *A TRS R is called* shallow *(respectively* flat*), if all its rewrite rules have the form $f(t_1, \ldots, t_n) \to g(s_1, \ldots, s_m)$ or $x \to g(s_1, \ldots, s_m)$ or $f(t_1, \ldots, t_n) \to x$ where every t_i and s_i is either a variable of $\mathcal{X}$ or a ground term of $\mathcal{T}(\Sigma)$ (respectively a variable of $\mathcal{X}$ or a symbol of Σ_0), and where $x \in \mathcal{X}$, and n, m can be 0 (if f or g have arity 0).*

We are interested in the following decision problems:

(ground) reachability. Given a TRS R on a signature Σ and two (ground) terms $s, t \in \mathcal{T}(\Sigma, \mathcal{X})$, do we have a reduction $s \xrightarrow[R]{*} t$?
(ground) joinability. Given a TRS R on Σ and two (ground) terms $s, t \in \mathcal{T}(\Sigma, \mathcal{X})$, does there exists $v \in \mathcal{T}(\Sigma, \mathcal{X})$ such that $s \xrightarrow[R]{*} v \xleftarrow[R]{*} t$?
confluence. Given a TRS R on Σ, do we have: for all $s, t \in \mathcal{T}(\Sigma, \mathcal{X})$ such that $s \xleftarrow[R]{*} u \xrightarrow[R]{*} t$ for some $u \in \mathcal{T}(\Sigma, \mathcal{X})$, does there exists $v \in \mathcal{T}(\Sigma, \mathcal{X})$ such that $s \xrightarrow[R]{*} v \xleftarrow[R]{*} t$?

We shall show below that the ground reachability, ground joinability and confluence problems are undecidable for flat TRS, by reduction of the Post correspondence problem.

[1] Reachability, joinability and confluence are shown NP-hard in [6].

2 Post Correspondence Problem, coding and coloring

We consider an instance of the Post Correspondence Problem (PCP) given by a finite set of pairs of words:

$$\text{PCP} := \{(u_i, v_i) \mid u_i, v_i \in \{a,b\}^*, 1 \leq i \leq N\} \tag{1}$$

The following problem is undecidable: [9]:

> Does there exist a finite sequence $(i_j)_{0\leq j\leq k}$ with $1 \leq i_0, \ldots, i_k \leq N$, such that $u_{i_0}u_{i_1}\ldots u_{i_k} = v_{i_0}v_{i_1}\ldots v_{i_k}$?

We shall represent the hypothetical solutions of PCP by ground terms from the sets described in Section 2.1, and provide in Sections 3, 4, and 5 some reductions to the reachability, joinability and confluence decision. The ingredients for the construction of the TRSs used in the reductions are two automata (Section 2.2), four TRSs (beginning of Section 3) and some coloring (Section 2.3).

2.1 Product and string terms

Let $_$ be a new symbol. We shall use a product operator $\otimes$ which associate to two words of $\{a,b\}^*$ a word of $\{a,b,_\}^{2^*}$ as follows:

$$c_1\ldots c_n \otimes c'_1\ldots c'_m := \langle c_1, c'_1\rangle \ldots \langle c_k, c'_k\rangle$$

where $c_1, \ldots, c_n, c'_1, \ldots, c'_m \in \{a,b\}$, $k = \max(n,m)$, and for all i with $n < i \leq k$, if any, (resp. all j with $m < j \leq k$), $c_i = _$ (resp. $c'_j = _$).

Example 2 $a \otimes bab = \langle a,b\rangle\langle _,a\rangle\langle _,b\rangle$.

Let us consider the signature $\Gamma := \{a, b, \varepsilon\}$, where a, b and ε have the respective arities 1,1 and 0.

We write: $\Gamma__ := \Gamma \uplus \{_\}$, where $_$ has arity 0 in $\Gamma__$, and $\Delta := \{a,b,_\}^2 \cup \{\varepsilon\}$, where ε has arity 0 in Δ, and every other symbols have arity 1 in Δ.

Remark 3 *We make no distinctions below between a word $c_1\ldots c_n \in \{a,b\}^*$ (resp. $d_1\ldots d_n \in \{a,b,_\}^{2^*}$) and the ground term $c_1(\ldots c_n(\varepsilon)) \in \mathcal{T}(\Gamma)$ (resp. $d_1(\ldots d_n(\varepsilon)) \in \mathcal{T}(\Delta)$).*

In this manner, the operator $\otimes$ is extended to $\mathcal{T}(\Gamma) \times \mathcal{T}(\Gamma) \to \mathcal{T}(\Delta)$.

2.2 Automata associated to PCP

Let A be a finite automaton computing on $\{a, b, _\}^{2^*}$, recognizing the set: $L(A) = \{u_i \otimes v_i \mid 1 \leq i \leq N\}^*$. We denote Q_A the set of states of A, and q_A its initial state.

Let B be a automaton with state set Q_B and initial state q_B recognizing $L(B) = \{a, b\}^*$.

Following Remark 3, we shall consider $L(A)$ and $L(B)$ as subsets of, respectively, $\mathcal{T}(\Delta)$ and $\mathcal{T}(\Gamma)$.

We associate to A and B two ground TRS T_A and T_B on the respective signatures $\Delta \uplus Q_A$ and $\Gamma \uplus Q_B$, where the states symbols of Q_A and Q_B have arity 0, as follows:

$$\begin{aligned} T_A := &\{q \to d(q') \mid q, q' \in Q_A,\ d \in \Delta,\ q \xrightarrow{d} q' \text{ is a transition of } A\} \\ &\cup \{q \to q' \mid q, q' \in Q_A,\ q \to q' \text{ is an epsilon-transition of } A\} \\ &\cup \{q \to \varepsilon \mid q \in Q_A \text{ is a final state of } A\} \qquad (2) \\ T_B := &\{q \to c(q') \mid q, q' \in Q_B,\ c \in \Gamma,\ q \xrightarrow{c} q' \text{ is a transition of } B\} \\ &\cup \{q \to q' \mid q, q' \in Q_B,\ q \to q' \text{ is an epsilon-transition of } B\} \\ &\cup \{q \to \varepsilon \mid q \in Q_B \text{ is a final state of } B\} \qquad (3) \end{aligned}$$

2.3 Coloring terms and TRS

We assume given 19 disjoint copies of the above signatures, colored with color i for $0 \leq i \leq 18$: $\Gamma_^{(i)} := \{c^{(i)} \mid c \in \Gamma_\}$, $Q_A^{(i)} := \{q^{(i)} \mid q \in Q_A\}$, $Q_B^{(i)} := \{q^{(i)} \mid q \in Q_B\}$, $\Delta^{(i)} := \{\langle c^{(i)}, c'^{(i)} \rangle \mid \langle c, c' \rangle \in \Delta\}$.

Let Θ be the following signature $\Theta := \Gamma_ \cup \Delta \cup Q_A \cup Q_B$, where the symbols of $\Gamma_$ and Δ keep their respective arities in Θ and the symbols of Q_A and Q_B have arity 0 in Θ, and let and $\Theta^{(i)}$ $(0 \leq i \leq 18)$ be the colored copies of Θ, $\Theta^{(i)} := \Gamma_^{(i)} \cup \Delta^{(i)} \cup Q_A^{(i)} \cup Q_B^{(i)}$.

For $0 \leq i \leq 18$, the i-coloring $t^{(i)} \in \mathcal{T}(\Theta^{(i)}, \mathcal{X})$ of a term of $t \in \mathcal{T}(\Theta, \mathcal{X})$ is recursively defined by: $f(t)^{(i)} := f^{(i)}(t^{(i)})$, and $x^{(i)} := x$ for all $x \in \mathcal{X}$.

Given a set $U \subseteq \mathcal{T}(\Theta)$, we write $U^{(i)} := \{t^{(i)} \mid t \in U\}$, and given a TRS R on Θ, we let $R^{(i,j)} := \{l^{(i)} \to r^{(j)} \mid l \to r \in R\}$ and $R^{(i)} := R^{(i,i)}$.

3 Reduction of PCP to reachability for flat TRS

We associate a TRS R_1 to the above problem PCP in (8), see also Figure 1. Its definition refers to the following two trivial and two projections TRS:

$$S := \{c(x) \to c(x) \mid c \in \Gamma_1\} \cup \{\varepsilon \to \varepsilon\} \tag{4}$$
$$P := \{d(x) \to d(x) \mid d \in \Delta_1\} \cup \{\varepsilon \to \varepsilon\} \tag{5}$$
$$\begin{aligned}\Pi_1 := &\{\langle c, c'\rangle(x) \to c(x) \mid c \in \Gamma_1, c' \in \Gamma_{_}\} \\ &\cup \{\langle _, c'\rangle(x) \to x \mid c' \in \Gamma_1\} \cup \{\varepsilon \to \varepsilon\}\end{aligned} \tag{6}$$
$$\begin{aligned}\Pi_2 := &\{\langle c, c'\rangle(x) \to c'(x) \mid c \in \Gamma_{_}, c' \in \Gamma_1\} \\ &\cup \{\langle c, _\rangle(x) \to x \mid c \in \Gamma_1\} \cup \cup\{\varepsilon \to \varepsilon\}\end{aligned} \tag{7}$$

The identity TRS S and P shall of course be used only in their colored form $S^{(i,j)}$ and $P^{(i,j)}$.

Example 4 *Assume that $u_1 = a$, $v_1 = bab$, $u_2 = ab$, $v_2 = b$. The term $(u_1 \otimes v_1)(u_2 \otimes v_2) \in L(A)$ is reduced as follows by the left projection Π_1 is, with an outermost strategy:*

$$\begin{aligned}&(u_1 \otimes v_1)(u_2 \otimes v_2) = \langle a, b\rangle(\langle _, a\rangle(\langle _, b\rangle(\langle a, b\rangle(\langle b, _\rangle(\varepsilon))))) \xrightarrow[\Pi_1]{} \\ &a(\langle _, a\rangle(\langle _, b\rangle(\langle a, b\rangle(\langle b, _\rangle(\varepsilon))))) \xrightarrow[\Pi_1]{} a(\langle _, b\rangle(\langle a, b\rangle(\langle b, _\rangle(\varepsilon)))) \xrightarrow[\Pi_1]{} \\ &a(\langle a, b\rangle(\langle b, _\rangle(\varepsilon))) \xrightarrow[\Pi_1]{} a(a(\langle b, _\rangle(\varepsilon))) \xrightarrow[\Pi_1]{} a(a(b(\varepsilon)))\end{aligned}$$

The TRS R_1 is defined on an extended signature: $\Xi = \bigcup_{i=0}^{i=18} \Theta^{(i)} \uplus \{f, g, 0, 1\}$ where $f, g, 0, 1$ are new function symbols of respective arities 8, 8, 0, 0 in Ξ.

$$R_1 := R_0 \cup \left\{ \begin{array}{l} 0 \to f(q_A^{(3)}, q_A^{(4)}, q_A^{(5)}, q_B^{(13)}, q_B^{(14)}, q_A^{(6)}, q_B^{(15)}, q_B^{(16)}), \\ f(x_1, x_2, x_1, y_{11}, y_{12}, x_2, y_{11}, y_{12}) \to g(x_1, x_2, x_1, y_{11}, y_{12}, x_2, y_{11}, y_{12}), \\ g(x_0, x_0, y_{17}, y_{17}, y_{18}, y_{18}, y_{10}, y_{10}) \to 1 \end{array} \right\}$$
$$\begin{aligned}R_0 := &T_A^{(3)} \cup T_A^{(4)} \cup T_A^{(5)} \cup T_A^{(6)} \cup T_B^{(13)} \cup T_B^{(14)} \cup T_B^{(15)} \cup T_B^{(16)} \cup \\ &P^{(3,1)} \cup P^{(4,2)} \cup P^{(5,1)} \cup S^{(13,11)} \cup S^{(14,12)} \cup P^{(6,2)} \cup S^{(15,11)} \cup S^{(16,12)} \cup \\ &P^{(1,0)} \cup P^{(2,0)} \cup \Pi_1^{(1,17)} \cup S^{(11,17)} \cup S^{(12,18)} \cup \Pi_2^{(2,18)} \cup S^{(11,10)} \cup S^{(12,10)}\end{aligned} \tag{8}$$

Note that R_1 is a flat TRS.

Definition 5 *A* 01-derivation witness *for R_1 is a tuple $(s_0, s_1, s_2, t_{10}, t_{11}, t_{12}, t_{17}, t_{18})$ of terms of $\mathcal{T}(\Xi, \mathcal{X})$ such that: $f(q_A^{(3)}, q_A^{(4)}, q_A^{(5)}, q_B^{(13)}, q_B^{(14)}, q_A^{(6)}, q_B^{(15)}, q_B^{(16)}) \xrightarrow[R_1]{*}$ $f(s_1, s_2, s_1, t_{11}, t_{12}, s_2, t_{11}, t_{12}) \xrightarrow[R_1]{} g(s_1, s_2, s_1, t_{11}, t_{12}, s_2, t_{11}, t_{12}) \xrightarrow[R_1]{*}$ $g(s_0, s_0, t_{17}, t_{17}, t_{18}, t_{18}, t_{10}, t_{10})$.*

$$
\begin{array}{llllllllll}
0 \to f(& q_A^{(3)}, & q_A^{(4)}, & q_A^{(5)}, & q_B^{(13)}, & q_B^{(14)}, & q_A^{(6)}, & q_B^{(15)}, & q_B^{(16)}) & \\
& T_A^{(3)} & T_A^{(4)} & T_A^{(5)} & T_B^{(13)} & T_B^{(14)} & T_A^{(6)} & T_B^{(15)} & T_B^{(16)} & \\
& P^{(3,1)} & P^{(4,2)} & P^{(5,1)} & S^{(13,11)} & S^{(14,12)} & P^{(6,2)} & S^{(15,11)} & S^{(16,12)} & \\
f(& x_1, & x_2, & x_1, & y_{11}, & y_{12}, & x_2, & y_{11}, & y_{12}) & \\
\downarrow & & & & & & & & & \\
g(& x_1, & x_2, & x_1, & y_{11}, & y_{12}, & x_2, & y_{11}, & y_{12}) & \\
& P^{(1,0)} & P^{(2,0)} & \Pi_1^{(1,17)} & S^{(11,17)} & S^{(12,18)} & \Pi_2^{(2,18)} & S^{(11,10)} & S^{(12,10)} & \\
g(& x_0, & x_0, & y_{17}, & y_{17}, & y_{18}, & y_{18}, & y_{10}, & y_{10}) & \to 1
\end{array}
$$

Fig. 1. The TRS R_1. The placement of the rules illustrates the equivalence between the existence of a solution to PCP and the existence of a reduction $0 \xrightarrow[R_1]{*} 1$. A solution is represented by a term $s \in L(A)$ such that $s \xrightarrow[\Pi_1]{*} t \xleftarrow[\Pi_2]{*} s$ for some $t \in L(B)$. The terms s and t are duplicated in the reduction (with different colors), they correspond to the variables x_i and y_j respectively. In the reduction, the rules of the top part (above $f(\ldots) \to g(\ldots)$) ensures that the (instances of) x_i and y_j belong respectively to $L(A)$ and $L(B)$ and the rules of the bottom part ensure the above relation between s and t, namely x_1 and x_2 are the same term x_0, modulo coloring, the projection with Π_1 of x_1 is y_{17}, the projection with Π_2 of x_2 is y_{18}, and y_{17} and y_{18} are the same term y_{10} modulo coloring.

Lemma 6 $0 \xrightarrow[R_1]{*} 1$ *iff there exists a 01-derivation witness for* R_1.

Proof: The *if* direction is immediate.

For the *only if* direction, let us remark that, by construction of R_1, the second and last-but-one terms of the derivation $0 \xrightarrow[R_1]{*} 1$ are necessarily respectively the first and last terms of the derivation in Definition 5, for some terms s_0, t_{17}, t_{18}, t_{10}.

Moreover, a case analysis on the rules of R_1 also shows that the derivation $f(q_A^{(3)}, q_A^{(4)}, q_A^{(5)}, q_B^{(13)}, q_B^{(14)}, q_A^{(6)}, q_B^{(15)}, q_B^{(16)}) \xrightarrow[R_1]{*} g(s_0, s_0, t_{17}, t_{17}, t_{18}, t_{18}, t_{10}, t_{10})$ has to involve the rule $f(x_1, x_2, x_1, y_{11}, y_{12}, x_2, y_{11}, y_{12}) \to g(x_1, x_2, x_1, y_{11}, y_{12}, x_2, y_{11}, y_{12})$ at the root position. This implies the existence of a 01-derivation witness. □

Lemma 7 *Every 01-derivation witness for* R_1 $w = (s_0, s_1, s_2, t_{10}, t_{11}, t_{12}, t_{17}, t_{18})$ *is such that:*

(1) $s_0 \in L(A)^{(0)}$, $s_1 \in L(A)^{(1)}$, $s_2 \in L(A)^{(2)}$, $t_{10} \in L(B)^{(10)}$, $t_{11} \in L(B)^{(11)}$, $t_{12} \in L(B)^{(12)}$, $t_{17} \in L(B)^{(17)}$ *and* $t_{18} \in L(B)^{(18)}$,

(2) $s_1 \xrightarrow[P^{(1,0)}]{*} s_0 \xleftarrow[P^{(2,0)}]{*} s_2$, *and* $t_{11} \xrightarrow[S^{(11,10)}]{*} t_{10} \xleftarrow[S^{(12,10)}]{*} t_{12}$,
(3) $s_1 \xrightarrow[\Pi_1^{(1,17)}]{*} t_{17} \xleftarrow[S^{(11,17)}]{*} t_{11}$, *and* $t_{12} \xrightarrow[S^{(12,18)}]{*} t_{18} \xleftarrow[\Pi_2^{(2,18)}]{*} s_2$.

Proof:

We can show (e.g. by induction on the length of the derivation) that every term t such that $f(q_A^{(3)}, q_A^{(4)}, q_A^{(5)}, q_B^{(13)}, q_B^{(14)}, q_A^{(6)}, q_B^{(15)}, q_B^{(16)}) \xrightarrow[R_1]{*} t$ verifies either $t = 1$ or t contains exactly one occurrence of f or g at root position and does not contain any occurrence of 0 or 1. Hence, all the reductions in $f(q_A^{(3)}, q_A^{(4)}, q_A^{(5)}, q_B^{(13)}, q_B^{(14)}, q_A^{(6)}, q_B^{(15)}, q_B^{(16)}) \xrightarrow[R_1]{*} f(s_1, s_2, s_1, t_{11}, t_{12}, s_2, t_{11}, t_{12})$ of Definition 5 and in $g(s_1, s_2, s_1, t_{11}, t_{12}, s_2, t_{11}, t_{12}) \xrightarrow[R_1]{*} g(s_0, s_0, t_{17}, t_{17}, t_{18}, t_{18}, t_{10}, t_{10})$ involve only rules of R_0 applied at non root positions.

Therefore, we have on one hand (see also Figure 1) $q_A^{(3)} \xrightarrow[R_0]{*} s_1$, $q_A^{(4)} \xrightarrow[R_0]{*} s_2$, $q_A^{(5)} \xrightarrow[R_0]{*} s_1$, $q_B^{(13)} \xrightarrow[R_0]{*} t_{11}$, $q_B^{(14)} \xrightarrow[R_0]{*} t_{12}$, $q_A^{(6)} \xrightarrow[R_0]{*} s_2$, $q_B^{(15)} \xrightarrow[R_0]{*} t_{11}$, $q_B^{(16)} \xrightarrow[R_0]{*} t_{12}$, and on the other hand $s_1 \xrightarrow[R_0]{*} s_0 \xleftarrow[R_0]{*} s_2$, $s_1 \xrightarrow[R_0]{*} t_{17} \xleftarrow[R_0]{*} t_{11}$, $t_{12} \xrightarrow[R_0]{*} t_{18} \xleftarrow[R_0]{*} s_2$, $t_{11} \xrightarrow[R_0]{*} t_{10} \xleftarrow[R_0]{*} t_{12}$.

We shall now show that the use of colors in the construction of R_0 implies (1)–(3) for w, by reducing incrementally the possible domain of each component of w.

- $q_A^{(3)} \xrightarrow[R_0]{*} s_1 \xleftarrow[R_0]{*} q_A^{(5)}$. Because of the choice of the colors 3 and 5, the left derivation can only involve rules of the sub systems $T_A^{(3)}$, $P^{(3,1)}$, $P^{(1,0)}$, $\Pi_1^{(1,17)}$, and the right derivation can only involve rules of the sub systems $T_A^{(5)}$, $P^{(5,1)}$, $P^{(1,0)}$, $\Pi_1^{(1,17)}$. Hence, $s_1 \in \mathcal{T}(\Delta^{(0)} \cup \Delta^{(1)} \cup \Gamma^{(17)})$.
- $q_A^{(4)} \xrightarrow[R_0]{*} s_2 \xleftarrow[R_0]{*} q_A^{(6)}$ implies similarly that $s_2 \in \mathcal{T}(\Delta^{(0)} \cup \Delta^{(2)} \cup \Gamma^{(18)})$.
- $s_1 \xrightarrow[R_0]{*} s_0 \xleftarrow[R_0]{*} s_2$ implies that $s_0 \in \mathcal{T}(\Delta^{(0)})$, $s_1 \in \mathcal{T}(\Delta^{(0)} \cup \Delta^{(1)})$ and $s_2 \in \mathcal{T}(\Delta^{(0)} \cup \Delta^{(2)})$.
- $q_B^{(13)} \xrightarrow[R_0]{*} t_{11} \xleftarrow[R_0]{*} q_B^{(15)}$ implies that $t_{11} \in \mathcal{T}(\Gamma^{(10)} \cup \Gamma^{(11)} \cup \Gamma^{(17)})$.
- $q_B^{(14)} \xrightarrow[R_0]{*} t_{12} \xleftarrow[R_0]{*} q_B^{(16)}$ implies that $t_{12} \in \mathcal{T}(\Gamma^{(10)} \cup \Gamma^{(12)} \cup \Gamma^{(18)})$.
- $t_{11} \xrightarrow[R_0]{*} t_{10} \xleftarrow[R_0]{*} t_{12}$ implies that $t_{10} \in \mathcal{T}(\Gamma^{(10)})$, $t_{11} \in \mathcal{T}(\Gamma^{(10)} \cup \Gamma^{(11)})$ and $t_{12} \in \mathcal{T}(\Gamma^{(10)} \cup \Gamma^{(12)})$.
- $s_1 \xrightarrow[R_0]{*} t_{17} \xleftarrow[R_0]{*} t_{11}$ implies that $s_1 \in \mathcal{T}(\Delta^{(1)})$, $t_{11} \in \mathcal{T}(\Gamma^{(11)})$ and $t_{17} \in \mathcal{T}(\Gamma^{(17)})$.
- $t_{12} \xrightarrow[R_0]{*} t_{18} \xleftarrow[R_0]{*} s_2$ implies that $s_2 \in \mathcal{T}(\Delta^{(2)})$, $t_{12} \in \mathcal{T}(\Gamma^{(12)})$ and $t_{18} \in \mathcal{T}(\Gamma^{(18)})$.

Altogether, we have shown (1). The conditions (2) and (3) follow then from the above reductions and the colors of the components of w. □

Lemma 8 *There exists a 01-derivation witness for* R_1 *iff there exists a solution for* PCP.

Proof: For the *if* direction, assume that the sequence $(i_j)_{0\leq j\leq k}$ is a solution of PCP, and let $s := (u_{i_0} \otimes v_{i_0})(u_{i_1} \otimes v_{i_1}) \dots (u_{i_k} \otimes v_{i_k})$ and $t := u_{i_0}u_{i_1} \dots u_{i_k}$. By construction of R_1, and because $t = v_{i_0}v_{i_1} \dots v_{i_k}$, the tuple $(s^{(0)}, s^{(1)}, s^{(2)}, t^{(10)}, t^{(11)}, t^{(12)}, t^{(17)}, t^{(18)})$ is a 01-derivation witness for R_1.

For the *only if* direction, let $(s_0, s_1, s_2, t_{10}, t_{11}, t_{12}, t_{17}, t_{18})$ be a 01-derivation witness for R_1. The conditions $s_0 \in L(A)^{(0)}$, $s_1 \in L(A)^{(1)}$, $s_2 \in L(A)^{(2)}$ of (1) in Lemma 7, and $s_1 \xrightarrow[P^{(1,0)}]{*} s_0 \xleftarrow[P^{(2,0)}]{*} s_2$ of (2) in Lemma 7 ensure that there exist a sequence $(i_j)_{0\leq j\leq k}$ such that, for each $\ell = 0, 1, 2$:

$$s_\ell = (u_{i_0}^{(\ell)} \otimes v_{i_0}^{(\ell)}) \cdots (u_{i_k}^{(\ell)} \otimes v_{i_k}^{(\ell)})$$

We can show that The other conditions in (1)–(3) in Lemma 7 imply that $(i_j)_{0\leq j\leq k}$ is a solution of PCP.

Indeed, $s_1 \xrightarrow[\Pi_1^{(1,17)}]{*} t_{17}$ and $t_{17} \in L(B)^{(17)})$ imply that $t_{17} = u_{i_0} \dots u_{i_k}(\varepsilon)^{(17)}$.

Also, $s_2 \xrightarrow[\Pi_2^{(2,18)}]{*} t_{18}$ and $t_{18} \in L(B)^{(18)})$ imply that $t_{18} = v_{i_0} \dots v_{i_k}(\varepsilon)^{(18)}$.

Moreover, $t_{11} \xrightarrow[S^{(11,17)}]{*} t_{17}$, $t_{12} \xrightarrow[S^{(12,18)}]{*} t_{18}$, and $t_{11} \in L(B)^{(11)}$, $t_{12} \in L(B)^{(12)}$, imply that $t_{11} = u_{i_0} \dots u_{i_k}(\varepsilon)^{(11)}$ and $t_{12} = v_{i_0} \dots v_{i_k}(\varepsilon)^{(12)}$.

Finally, $t_{11} \xrightarrow[S^{(11,10)}]{*} t_{10} \xleftarrow[S^{(12,10)}]{*} t_{12}$, permits us to conclude that $u_{i_0} \dots u_{i_k}(\varepsilon) = v_{i_0} \dots v_{i_k}(\varepsilon)$. □

Lemmas 6, 7 and 8 establish a reduction of the undecidable PCP into the reachability problem for $(R_1, 0, 1)$. Hence we can conclude with the following theorem.

Theorem 9 *The ground reachability problem is undecidable for flat TRS.*

4 Reduction of PCP to joinability for flat TRS

The undecidability for the joinability follows from a reduction presented in [7] (we can also observe that the joinability problem for R_1, 0 and 1 is equivalent to the reachability problem for R_1, 0 and 1).

Corollary 10 *The ground joinability problem is undecidable for flat TRS.*

5 Reduction of PCP to confluence for flat TRS

We shall modify the TRS R_1 built for the proof of Theorem 9 in order to reduce PCP to confluence [2] . More precisely, we shall construct a TRS R_2 such that $0 \xrightarrow[R_1]{*} 1$ iff $R_1 \cup R_2$ is confluent.

The TRS R_2 is defined on the extended signature: $\Xi' = \Xi \cup \{2\}$, where 2 has arity 0 in Ξ'.

$$\begin{aligned} R_2 := \{2 \to 0, 2 \to 1\} \cup \{c \to 0 \mid c \in \Xi_0 \setminus \{0,1\}\} \cup \{d(x) \to 0 \mid d \in \Xi_1\} \cup \\ \{d(1) \to 1 \mid d \in \Xi_1\} \cup \{f(z_1, \ldots, z_8) \to 1, g(z_1, \ldots, z_8) \to 1 \mid \\ \text{one of the } z_i \text{ is } 1, \text{the others are distinct variables}\} \end{aligned} \tag{9}$$

We recall that Ξ_0 and Ξ_1 denote the set of symbols of Ξ of arity respectively 0 and 1. Note that R_2 is flat.

Lemma 11 *$R_1 \cup R_2$ is confluent iff $0 \xrightarrow[R_1]{*} 1$.*

Proof: For the *only if* direction, assume that $0 \not\xrightarrow[R_1]{*} 1$. It means that 0 and 1 are not joinable by R_1 (since 1 is in normal form for $R_1 \cup R_2$) and hence also not joinable by $R_1 \cup R_2$. Hence $R_1 \cup R_2$ is not confluent because of the peak $0 \xleftarrow[R_2]{} 2 \xrightarrow[R_2]{} 1$.

For the *if* direction, assume that $0 \xrightarrow[R_1]{*} 1$, and let $R_3 := (R_1 \cup R_2) \setminus R_4$, where R_4 contains the rules of $T_A^{(3)}$, $T_A^{(4)}$, $T_A^{(5)}$, $T_A^{(6)}$, $T_B^{(13)}$, $T_B^{(14)}$, $T_B^{(15)}$, $T_B^{(16)}$ and $0 \to f(q_A^{(3)}, q_A^{(4)}, q_A^{(5)}, q_B^{(13)}, q_B^{(14)}, q_A^{(6)}, q_B^{(15)}, q_B^{(16)})$, $2 \to 0$, $2 \to 1$.

Let u, t, s be some terms such that $s \xleftarrow[R_1 \cup R_2]{*} u \xrightarrow[R_1 \cup R_2]{*} t$. Suppose that the reduction $u \xrightarrow[R_1 \cup R_2]{*} s$ uses some rules of R_4. It means that some term in this reduction contains a constant symbol of Ξ_0. We can show (by induction on the reduction length) that if $s_1 \xrightarrow[R_1 \cup R_2]{*} s_2$ then s_1 contains a constant of Ξ_0 iff s_2 also contains a constant of Ξ_0. Hence in our case, u, s and t contain at least a constant of Ξ_0. It implies that $s \xrightarrow[R_2]{*} 1 \xleftarrow[R_2]{*} t$ (because $0 \xrightarrow[R_1]{*} 1$).

Hence, it is sufficient to show that R_3 is confluent in order to show that $R_1 \cup R_2$ is confluent. We can show that R_3 is terminating, by exhibiting a reduction ordering compatible with R_3, for instance, a multiset path ordering [10] based on a well founded precedence $\succ$ on Ξ such that:

$d \succ c$ for all $d \in \Delta^{(i)}$, $c \in \Gamma_{_}^{(j)}$, $0 \leq i, j \leq 18$,

[2] A similar technique is used in [6] to show the NP-hardness of confluence for shallow TRS.

$c^{(i)} \succ c^{(j)}$ for all $c \in \Gamma$, and $(i,j) = (13,11),\ (14,12),\ (15,11),\ (16,12),\ (11,17),\ (12,18),\ (11,10),\ (12,10)$,
$d^{(i)} \succ d^{(j)}$ for all $d \in \Delta$, and $(i,j) = (3,1),\ (4,2),\ (5,1),\ (6,2),\ (1,0),\ (2,0)$,
$\varepsilon^{(1)} \succ \varepsilon^{(17)}$, $\varepsilon^{(2)} \succ \varepsilon^{(18)}$,
$c \succ 0$ for all $c \in \Xi_0 \cup \Xi_1$, $f \succ g$, $g \succ 1$.

Using Newman's lemma, we show the confluence of R_3 by observing that all its critical pairs can be joined by R_2. We detail below all the cases of critical pairs.

Each of the following critical peaks of the form $s \xleftarrow[R_3]{} \ell \xrightarrow[R_3]{} t$ can be joined in zero or one step with a rule of R_2 (i.e. $s \xrightarrow[R_2]{*} t$):

- $\varepsilon^{(j)} \xleftarrow[R]{} \varepsilon^{(i)} \xrightarrow[R_2]{} 0$, where $R = S^{(i,j)} \cup P^{(i,j)} \cup \Pi_1^{(i,j)} \cup \Pi_2^{(i,j)}$,
- $d'^{(j)}(x) \xleftarrow[R]{} d^{(i)}(x) \xrightarrow[R_2]{} 0$, where $R = S^{(i,j)} \cup P^{(i,j)} \cup \Pi_1^{(i,j)} \cup \Pi_2^{(i,j)}$ and $d, d' \in \Xi_1$,
- $d'^{(j)}(1) \xleftarrow[R]{} d^{(i)}(1) \xrightarrow[R_2]{} 1$ where $R = S^{(i,j)} \cup P^{(i,j)} \cup \Pi_1^{(i,j)} \cup \Pi_2^{(i,j)}$ and $d, d' \in \Xi_1$,
- $g(1, z_2, 1, z_4, z_5, z_2, z_4, z_5) \xleftarrow[R_1]{} f(1, z_2, \ldots, z_8) \xrightarrow[R_2]{} 1$, and similarly for any rule $f(z_1, \ldots, z_8) \to 1 \in R_2$ such that one $z_i = 1$,
- $1 \xleftarrow[R_1]{} g(1, z_2, \ldots, z_8) \xrightarrow[R_2]{} 1$, and similarly for any rule $g(z_1, \ldots, z_8) \to 1 \in R_2$ such that one $z_i = 1$.

Each of the following critical pairs of the form $s \xleftarrow[R_3]{} \ell \xrightarrow[R_3]{} t$ can be joined in two steps with R_2 by $s \xrightarrow[R_2]{} 0 \xleftarrow[R_2]{} t$:

- $\varepsilon^{(j)} \xleftarrow[R]{} \varepsilon^{(i)} \xrightarrow[R']{} \varepsilon^{(j')}$, where $R = S^{(i,j)} \cup P^{(i,j)} \cup \Pi_1^{(i,j)} \cup \Pi_2^{(i,j)}$, $R' = S^{(i,j')} \cup P^{(i,j')} \cup \Pi_1^{(i,j')} \cup \Pi_2^{(i,j')}$,
- $c_1^{(j)}(x) \xleftarrow[R]{} {c_0}^{(i)}(x) \xrightarrow[R']{} {c_2}^{(j')}(x)$, where $R = S^{(i,j)} \cup P^{(i,j)} \cup \Pi_1^{(i,j)} \cup \Pi_2^{(i,j)}$, $R' = S^{(i,j')} \cup P^{(i,j')} \cup \Pi_1^{(i,j')} \cup \Pi_2^{(i,j')}$, and $c_0, c_1, c_2 \in \Xi_1$,
- $\langle c^{(0)}, c'^{(0)} \rangle(x) \xleftarrow[P^{(1,0)}]{} \langle c^{(1)}, c'^{(1)} \rangle(x) \xrightarrow[\Pi_1^{(1,17)}]{} c^{17}(x)$,
- $\langle c^{(0)}, c'^{(0)} \rangle(x) \xleftarrow[P^{(2,0)}]{} \langle c^{(2)}, c'^{(2)} \rangle(x) \xrightarrow[\Pi_2^{(2,18)}]{} c'^{18}(x)$.

□

By Lemmas 6, 7, 8 and 11, there is a reduction of PCP into the confluence for the flat TRS $R_1 \cup R_2$.

Theorem 12 *The confluence is undecidable for flat TRS.*

Conclusion

We have shown that the properties of reachability, joinability and confluence are undecidable for flat (and hence shallow) TRS. This is a big contrast with the shallow linear case, for which all these properties are known to be decidable in polynomial time [6].

One can note that all the known decidability results for confluence concern classes of linear TRS. Two subclasses of (non-linear) shallow TRS remain out of the scope of the reductions constructed here: the shallow right-ground TRS – reachability and joinability are decidable for right-ground rewrite systems [11], and, more generally, subclasses of shallow TRS with syntactic restriction on the relative occurrences of variables between left and right members of rules, e.g. shallow TRS such that a variable with more than one occurrence in the left member can not occur in the right member of a rewrite rule.

Acknowledgements

The author wishes to thank Ralf Treinen who has carefully read and commented a preliminary version of this paper, and Harald Ganzinger for his useful suggestions.

References

[1] M. Dauchet, T. Heuillard, P. Lescanne, S. Tison, Decidability of the confluence of finite ground term rewrite systems and of other related term rewrite systems, Information and Computation 88 (1990) 187–201.

[2] M. Oyamaguchi, The Church-Rosser property for ground term-rewriting systems is decidable, Theoretical Computer Science 49 (1) (1987) 43–79.

[3] A. Hayrapetyan, R. Verma, On the complexity of confluence for ground rewrite systems, in: Bar-Ilan International Symposium on the Foundations of Artificial Intelligence, 2002.

[4] H. Comon, G. Godoy, R. Nieuwenhuis, The confluence of ground term rewrite systems is decidable in polynomial time, in: 42nd Annual IEEE Symposium on Foundations of Computer Science, Las Vegas, Nevada, USA, 2001.

[5] A. Tiwari, Deciding confluence of certain term rewriting systems in polynomial time, in: G. Plotkin (Ed.), IEEE Symposium on Logic in Computer Science, IEEE Society, 2002, 447–456.

[6] G. Godoy, A. Tiwari, R. Verma, On the confluence of linear shallow term rewrite systems, in: H. Alt (Ed.), 20th Intl. Symposium on Theoretical Aspects of Computer Science, Lecture Notes in Computer Science, Springer, 2003, to appear.

[7] R. Verma, M. Rusinowitch, D. Lugiez, Algorithms and reductions for rewriting problems, Fundamenta Informaticae 43 (3) (2001) 257–276, also in Proc. of Intl Conference on Rewriting Techniques and Applications 1998.

[8] H. Ganzinger, F. Jacquemard, M. Veanes, Rigid reachability, in: Proc. Asian Computing Science Conference, Vol. 1538 of Lecture Notes in Computer Science, Springer-Verlag, Berlin, 1998.

[9] E. L. Post, A variant of a recursively unsovable problem, Bulletin of the AMS 52 (1946) 264–268.

[10] N. Dershowitz, Orderings for term rewriting systems, Theoretical Computer Science 17 (3) (1982) 279–301.

[11] M. Oyamaguchi, The reachability and joinability problems for right-ground term-rewriting systems, Journal of Information Processing 13 (3) (1990) 347–354.